

Basics of the theory of cyclic rook polynomials and cyclic permanents of rectangular matrices

A. M. Kamenetskii

Let A be a $k \times m$ matrix, $n \leq k$, and let $j^{(i)}, j_1^{(i)}, \dots, j_{p_i}^{(i)} \in N_m$, $1 \leq i \leq q$. Then, by definition, $A[N_n | j^{(i)} \oplus \sum_{l=1}^{p_i} \oplus j_l^{(i)}] = A[N_n | j^{(i)}] + \sum_{l=1}^{p_i} A[N_n | j_l^{(i)}]$ and $A[N_n | j^{(1)} \oplus \sum_{l=1}^{p_1} \oplus j_l^{(1)}, \dots, j^{(q)} \oplus \sum_{l=1}^{p_q} \oplus j_l^{(q)}]$ is a $n \times q$ matrix, whose i th column is $A[N_n | j^{(i)} \oplus \sum_{l=1}^{p_i} \oplus j_l^{(i)}]$.

Theorem 1 *Let $0 \leq r \leq n$, $n \geq 1$, $k \geq 1$, $t \geq 0$. Then*

$$\begin{aligned} \left(\sum_{i=0}^t a_i P_n^{-r+i} \right) \otimes J_k &= \left(\left(\sum_{i=0}^t a_i T_{n+t}^{(i)} \right) \otimes J_k \right) [N_{nk} | (k(r+1) \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv (r+1) \bmod n}} \oplus k(n+i))^{(k)}, \\ &\quad (k(r+2) \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv (r+2) \bmod n}} \oplus k(n+i))^{(k)}, \dots, (kn \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv n \bmod n}} \oplus k(n+i))^{(k)}, \\ &\quad (k \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv 1 \bmod n}} \oplus k(n+i))^{(k)}, (2k \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv 2 \bmod n}} \oplus k(n+i))^{(k)}, \dots, \\ &\quad (kr \oplus \sum_{\substack{1 \leq i \leq t \\ i \equiv r \bmod n}} \oplus k(n+i))^{(k)}]. \end{aligned}$$

An injective mapping φ of a set A induces the following bijective mapping φ^* of the set $\varphi(A) \setminus (\varphi(A) \cap A)$ onto $A \setminus (A \cap \varphi(A))$: if $j \in \varphi(A) \setminus (\varphi(A) \cap A)$, then $\varphi^*(j) = j_m$, where j_m is uniquely determined by the conditions $j_0 = j$ and $j_1, j_2, \dots, j_{m-1} \in \varphi(A)$, $j_l = \varphi^{-1}(j_{l-1})$, $1 \leq l \leq m$, $j_m \in A \setminus (\varphi(A) \cap A)$. Let $\bar{\alpha} = (x_1, \dots, x_n)$ be a sequence, $A \subseteq N_n$, $\varphi \in \text{Inj}(A, N_n)$. Denote by $\bar{\varphi}(\bar{\alpha})$ the sequence obtained from $\bar{\alpha}$ in the following way. The components x_j , $j \in \varphi(A)$, are deleted. And if $j \in \varphi(A) \setminus (\varphi(A) \cap A)$, then the deleted component x_j is replaced by $x_{\varphi^*(j)}$.

Theorem 2 (Expansion of the cyclic rook polynomial of a rectangular matrix along the last k rows) *Let $A = (a_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$ be an $m \times n$ matrix over a commutative ring with unity, $m \leq n$, $1 \leq k \leq m-1$. Then*

$$\begin{aligned} R(x; z; A) &= \sum_{S \subseteq N_m \setminus N_{m-k}, S \neq \emptyset} \sum_{\varphi \in \text{Inj}(S, N_n)} z^{|\varphi|} \left(\prod_{i \in S} a_{i, \varphi(i)} \right) R(x; z; A[N_{m-k} | \bar{\varphi}(N_n)] x^{|S|} \\ &\quad + R(x; z; A[N_{m-k} | N_n]). \end{aligned}$$

If $k > 1$, then such an expansion along arbitrary k rows, for an arbitrary matrix, does not exist. On the other hand, if $k = 1$, then such an expansion along arbitrary row exists.

Theorem 3 Let $A = (a_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$ and $m \leq n$, $1 \leq i \leq m$. Then

$$\begin{aligned} R(x; z; A) &= a_{i,i} x z R(x; z; A[N_m \setminus (i) | N_n \setminus (i)]) \\ &\quad + \sum_{j \in N_n \setminus (i)} a_{i,j} x R(x; z; A[N_m \setminus (i) | 1, 2, \dots, i-1, i+1, \dots, j-1, i, j+1, \dots, n]) \\ &\quad + R(x; z; A[N_m \setminus (i) | N_m \setminus (i), i, m+1, \dots, n]). \end{aligned}$$

Theorem 4 (Expansion of the cyclic rook polynomial along arbitrary k rows) Let $A = (a_{i,j})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$ $m \leq n$, $1 \leq k \leq m-1$, $\bar{\beta} \in Q_{k,m}$. Then

$$\text{per}(z; A) = \sum_{\varphi \in \text{Inj}(\{\bar{\beta}\}, N_n)} z^{|\varphi|} \left(\prod_{i \in \{\bar{\beta}\}} a_{i, \varphi(i)} \right) \text{per}(z; A[N_m \setminus \bar{\beta} | \bar{\varphi}(N_n)]).$$

Theorem 5 Let $A = (a_{i,j})$ and $B = (b_{i,j})$ be $m \times n$ matrices over a commutative ring with unity, $m \leq n$, $R(x; z; A) = \sum_{l=0}^m r_l(z; A) x^l$, $r_0(z; A) = 1$, $R(x; A) = \sum_{l=0}^m r_l(A) x^l$, $r_0(A) = 1$. Then

$$\begin{aligned} R(x; z; A + B) &= \sum_{s=0}^m \sum_{\bar{\alpha} \in Q_{s,m}} \sum_{\varphi \in \text{Inj}(\{\bar{\alpha}\}, N_n)} z^{|\varphi|} \left(\prod_{i \in \{\bar{\alpha}\}} a_{i, \varphi(i)} \right) x^s R(x; z; B[N_m \setminus \bar{\alpha} | \bar{\varphi}(N_n)]); \\ r_l(z; A + B) &= \sum_{s=0}^l \sum_{\bar{\alpha} \in Q_{s,m}} \sum_{\varphi \in \text{Inj}(\{\bar{\alpha}\}, N_n)} z^{|\varphi|} \left(\prod_{i \in \{\bar{\alpha}\}} a_{i, \varphi(i)} \right) r_{l-s}(z; B[N_m \setminus \bar{\alpha} | \bar{\varphi}(N_n)]); \\ \text{per}(z; A + B) &= \sum_{s=0}^m \sum_{\bar{\alpha} \in Q_{s,m}} \sum_{\varphi \in \text{Inj}(\{\bar{\alpha}\}, N_n)} z^{|\varphi|} \left(\prod_{i \in \{\bar{\alpha}\}} a_{i, \varphi(i)} \right) \text{per}(z; B[N_m \setminus \bar{\alpha} | \bar{\varphi}(N_n)]); \\ R(x; A + B) &= \sum_{s=0}^m \sum_{\substack{\bar{\alpha} \in Q_{s,m} \\ \bar{\beta} \in Q_{s,n}}} \text{per}(A[\bar{\alpha} | \bar{\beta}]) x^s R(x; B(\bar{\alpha} | \bar{\beta})); \\ r_l(A + B) &= \sum_{s=0}^l \sum_{\substack{\bar{\alpha} \in Q_{s,m} \\ \bar{\beta} \in Q_{s,n}}} \text{per}(A[\bar{\alpha} | \bar{\beta}]) r_{l-s}(B(\bar{\alpha} | \bar{\beta})); \\ \text{per}(A + B) &= \sum_{s=0}^m \sum_{\substack{\bar{\alpha} \in Q_{s,m} \\ \bar{\beta} \in Q_{s,n}}} \text{per}(A[\bar{\alpha} | \bar{\beta}]) \text{per}(B(\bar{\alpha} | \bar{\beta})). \end{aligned}$$

Theorem 6 Let A be an $m \times n$ matrix over a commutative ring with unity, $m \leq n$, $R(x; z; A) = \sum_{l=0}^m r_l(z; A) x^l$, $r_0(z; A) = 1$, $(z)^{(k)} = \sum_{i=0}^{k-1} (z+i)$, $k \geq 1$, $(z)^{(0)} = 1$. Then

$$r_l(z; y J_{m,n} - A) = \sum_{s=0}^l (-1)^s \binom{m-s}{l-s} r_s(z; A) (n-l+z)^{(l-s)} y^{l-s}, \quad 0 \leq l \leq m.$$

In particular,

$$\text{per}(z; yJ_{m,n} - A) = \sum_{s=0}^m (-1)^s r_s(z; A) (n - m + z)^{(m-s)} y^{m-s}.$$

Theorem 7

$$\text{per}(z; (a_0 I_n + a_1 P_n) \otimes J_k) = \sum_{s=0}^k \frac{(z)^{(s)}}{s!} [s! \binom{k}{s} a_0^{k-s} a_1^s (s + z)^{(k-s)}]^n$$

for all $n \geq 1$.

Theorem 8 Let $r \geq t + 1$. Then for all $n \geq 1$ we have

$$R(x; z; (\sum_{i=0}^t a_i P_n^{-r+i}) \otimes J_k) = \sum_{\substack{(l_1, \dots, l_t) \\ 0 \leq l_i \leq k, 1 \leq i \leq t}} \sum_M f(z; \begin{pmatrix} v_1^{(l_1)}, v_2^{(l_2)}, \dots, v_t^{(l_t)}, v_{t+1}^{(k)}, v_{t+2}^{(k)}, \dots, v_r^{(k)} \\ \bar{\gamma} \setminus (1^{(k-l_1)}, 2^{(k-l_2)}, \dots, t^{(k-l_t)}) \end{pmatrix}).$$

$$(K_{[k],t}^{(0,r)}(a_0 x, a_1 x, \dots, a_t x; z))^n [L(1^{(k-l_1)}, v_1^{(l_1)}, 2^{(k-l_2)}, v_2^{(l_2)}, \dots, t^{(k-l_t)}, v_t^{(l_t)}, v_{t+1}^{(k)}, v_{t+2}^{(k)}, \dots, v_r^{(k)}) | L(\bar{\gamma})],$$

where the summation is carried over the domain M defined by the conditions

$$\begin{aligned} \bar{\gamma} &\in \cup_{d=0}^{rk} G_{[k],d,t}^{(0,r)}, \\ \{\bar{\gamma}\} \setminus (\{\bar{\gamma}\} \cap \{w^{(rk)}\}) &= \{1^{(k-l_1)}, v_1^{(l_1)}, 2^{(k-l_2)}, v_2^{(l_2)}, \dots, t^{(k-l_t)}, v_t^{(l_t)}, v_{t+1}^{(k)}, v_{t+2}^{(k)}, \dots, v_r^{(k)}\}. \end{aligned}$$

Lemma 1 Let $\bar{\alpha} = (x_1, \dots, x_m)$ and $\bar{\gamma} = (y_1, \dots, y_n)$ be two sequences, $m \leq n$, $\{\bar{\alpha}\} \subseteq \{\bar{\gamma}\}$, $\{\bar{\alpha}\} = \{b_1^{(l_1)}, \dots, b_d^{(l_d)}\}$, $A_i = \{j \in N_m | x_j = b_i\}$, $B_i = \{j \in N_n | y_j = b_i\}$, $1 \leq i \leq d$. Then, for all i , $1 \leq i \leq d$, we have

$$f(z; \begin{pmatrix} \bar{\alpha} \\ \bar{\gamma} \end{pmatrix}) = \left(\binom{|B_i|}{|A_i|} (|A_i|!) \right)^{-1} \sum_{\varphi \in \text{Inj}(A_i, B_i)} z^{|\varphi|} f(z; \begin{pmatrix} \bar{\alpha} \setminus (b_i^{(l_i)}) \\ \bar{\varphi}(\bar{\gamma}) \end{pmatrix}).$$

The author expresses his deep gratitude to S. K.Lando for invaluable support and attention to the work.

References

- [1] A. M. Kamenetskii, Russian Mathematical Surveys, **60**:3 (2005), 177–178
 - [2] A. M. Kamenetskii, Russian Mathematical Surveys, **62**:6 (2007), 175–176
 - [3] A. M. Kamenetskii, Russian Mathematical Surveys, **63**:4 (2008), 187–188
- e-mail: `kamenetsky_a@list.ru`